



Bayer CropScience LP  
 P.O. Box 12014  
 2 T.W. Alexander Drive  
 Research Triangle Park, North Carolina 27709  
 1-866-99BAYER (1-866-992-2937)

ALIETTE® WDG Fungicide  
 For use as a Pea Seed Treatment for Control of Downy Mildew  
 EPA Reg. No. 264-516  
 EPA SLN. No. ID-030001

## *FIFRA 24(c) Special Local Need Label*

### FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

**THIS LABEL IS VALID UNTIL DECEMBER 31, 2017 OR UNTIL OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.**

#### DIRECTIONS FOR USE

##### MIXING INSTRUCTIONS

1. Fill the slurry tank with 1/4 - 1/2 of the required volume of water prior to the addition of ALIETTE® WDG.
2. Add ALIETTE® WDG slowly to the tank and agitate by hydraulic or mechanical means.

Note: Bayer CropScience has determined that phytotoxicity from solubilized copper may occur if products containing copper are tank-mixed with ALIETTE® WDG or if unbuffered ALIETTE® WDG is applied to foliage with copper residues.

Do not tank-mix with copper compounds. When applied prior to or after copper compounds, the pH of ALIETTE® WDG should be raised to 6.0 or above with the addition of an alkaline buffer such as Potassium Carbonate (3 lbs Potassium Carbonate to 5 lbs ALIETTE® WDG) or DiAmmonium Phosphate (5 lbs DiAmmonium Phosphate to 5 lbs ALIETTE® WDG).

3. Slowly add alkaline buffering agent (if needed).

CAUTION: When Potassium Carbonate is used as a buffer Carbon Dioxide (CO<sup>2</sup>) gas is released during the buffering process. The spray tank must NOT be sealed until the CO<sup>2</sup> release associated with the reaction subsides (approximately 5 minutes).

4. Continue to fill the tank with water and/or additional compatible seed treatments to the desired volume while agitating.
5. Continue agitation when applying.

Crop/Site:	Succulent, <b>shelled</b> pea varieties – Seed treatment use
Target Pest:	Downy Mildew ( <i>Peronospora viciae</i> )
Benefits:	Control of primary downy mildew Suppression of secondary downy mildew Effective on downy mildew strains resistant to metalaxyl and mefenoxam
Rate:	0.1875 lbs (3 oz) of ALIETTE® WDG per 100 lbs of seed
Method of Application:	Seed treatment: Treat peas using a slurry of fungicides at the rate of 7.7 fluid ounces of slurry per 100 pounds of seed. Maintain continuous agitation of slurry during application. Allow treated seeds to dry thoroughly before packaging or planting.
Use Restrictions:	Maximum of 1 application to pea seeds. For use by commercial seed treaters. Do not use treated seed for any other purpose than planting. An EPA-approved dye (e.g., 40 CFR 180.1001) which imparts an unnatural color to the seed, must be applied to treated seed because ALIETTE® WDG Fungicide is manufactured without a dye.
Labeling:	Federal law requires that bags containing treated seeds must be labeled with the following information: <b>“This seed has been treated with ALIETTE® WDG Fungicide. Do not use for feed, food, or oil purposes. Store away from feed and foodstuffs.”</b>

This product is toxic to aquatic and estuarine invertebrates. ALIETTE® WDG Fungicide should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may contact the National Marine Fisheries Service or US Fish & Wildlife Service for information on listed threatened or endangered species (e.g. Bull trout, Chinook salmon). Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

Aliette is a registered trademark of Bayer.

2/22/2013