

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.



FIFRA 24 (C)
Special Local Need Label
 Legume Vegetables (Succulent & Dried)

EPA Reg. No. 279-3302

EPA SLN. No. ID-150003

FIFRA Section 24 (c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF
IDAHO

This label is valid until December 31, 2018 or until otherwise amended, withdrawn, canceled, or suspended.

This label and the federal label for this product must be in the possession of the user at the time of pesticide application.
 Follow all applicable directions, restrictions, and precautions on this label and the main EPA-registered label.
 It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Crop	Pest		Rate of Application
SUCCULENT SHELLED PEAS AND BEANS Pre-Plant Incorporated (PPI)	Wireworm Grape colapsis Root maggot Armyworm spp.	Seed corn maggot White grubs Black cutworm	5.3 - 8.5 fluid oz/acre (0.062 - 0.1 lb ai/ acre)

DIRECTIONS FOR USE: Succulent shelled pea and beans subgroup.

FOR PRE-PLANT INCORPORATED APPLICATIONS (PPI) TO SUCCULENT SHELLED PEA AND BEANS

Bean (*Phaseolus*) (includes lima bean, green; broad bean, succulent); bean (*Vigna*) (includes blackeyed pea, cowpea, southern pea); pea (*Pisum*) (includes English pea, garden pea, green pea); pigeon pea.

Do not apply more than 0.2 pound active ingredient per acre per season to dry peas including pre-plant ppi, at-plant and foliar applications of other bifenthrin containing products.

Do not apply more than 0.1 pound active ingredient per acre per season as a pre-plant, PPI, or at-plant application.

When applied as directed, Capture LFR will provide control of the pests listed in the table above. For PPI treatments, the 5.3-8.5 oz/A rate must be used. Capture LFR can be tank mixed and applied with PPI herbicides. Incorporation of Capture LFR should no deeper than 3 inches. Incorporation should be close to the intended seed planting depth.

Crop	Pest	Rate of Application								
<p align="center">DRIED SHELLLED PEAS AND BEANS Pre-Plant Incorporated (PPI)</p>	<table border="0"> <tr> <td>Wireworm</td> <td>Seed corn maggot</td> </tr> <tr> <td>Grape colapsis</td> <td>White grubs</td> </tr> <tr> <td>Root maggot</td> <td>Black cutworm</td> </tr> <tr> <td>Armyworm spp.</td> <td></td> </tr> </table>	Wireworm	Seed corn maggot	Grape colapsis	White grubs	Root maggot	Black cutworm	Armyworm spp.		<p align="center">5.3 - 8.5 fluid oz/acre (0.062 - 0.1 lb ai/ acre)</p>
Wireworm	Seed corn maggot									
Grape colapsis	White grubs									
Root maggot	Black cutworm									
Armyworm spp.										

DIRECTIONS FOR USE: Dried shelled pea and bean (except soybean) subgroup.

FOR PRE-PLANT INCORPORATED APPLICATIONS (PPI) TO DRIED SHELLLED PEA AND BEANS.

Dried cultivars of bean (*Lupinus*) ; bean (*Phaseolus*) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); bean (*Vigna*) (includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); broad bean (dry); chickpea; guar; lablab bean; lentil; pea (*Pisum*) (includes field pea); pigeon pea.

When applied as directed, Capture LFR will provide control of the pests listed in the table above. For PPI treatments, the 5.3-8.5 oz/A rate must be used. Capture LFR can be tank mixed and applied with PPI herbicides. Incorporation of Capture LFR should no deeper than 3 inches. Incorporation should be close to the intended seed planting depth.

Restrictions:

Do not apply more than 0.2 pound active ingredient per acre per season to succulent peas and beans including pre, ppi, at-plant and foliar applications of other bifenthrin containing products.

Do not apply more than 0.3 pounds active ingredient per acre per season to dry beans including pre, ppi, at-plant and foliar applications of other bifenthrin containing products.

Do not apply more than 0.1 pound active ingredient per acre per season as a pre-plant, PPI, or at-plant application.

FMC is a trademark of FMC and Capture LFR Corporation. Capture LFR is a restricted use pesticide. All rights reserved.



FMC Corporation
Agricultural Solutions
1735 Market Street
Philadelphia, PA 19103

©All rights reserved Rev: 1-20-2015