



DuPont Crop Protection

SPECIAL LOCAL NEED 24(C)LABELING

DuPont™ ManKocide® fungicide/bactericide FOR USE ON CARROTS GROWN FOR SEED

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

DuPont™ ManKocide® fungicide/bactericide

EPA Reg. No. 352-690 EPA SLN No. ID-080012

FOR USE ON CARROTS GROWN FOR SEED TO PREVENT BACTERIAL BLIGHT

This label is valid until December 31, 2016, or until otherwise amended, withdrawn, canceled, or suspended.

PRECAUTIONS AND PPE REQUIREMENTS ON THE EPA-REGISTERED LABEL.

Table with 2 columns: Ingredient Name and Percentage. Includes Active Ingredients (Mancozeb, Manganese, Zinc, Ethylenebisdithiocarbamate ion, Copper Hydroxide) and Inert Ingredients.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Instructions for Control of Bacterial blight (Xanthomonas campestris)

Timing and Frequency of Application: Apply prior to conditions conducive to infection, such as precipitation, overhead irrigation, and frost and winds strong enough to create plant wounding via blown sand and dirt. Repeat applications on a 7-14 day interval depending on plant growth and disease pressure.

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

IMPORTANT

BEFORE USING MANKOCIDE®, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARD REQUIREMENTS,

Apply ManKocide® prior to establishment of infections. ManKocide® will not eradicate infections which are already established, but will provide partial protection from the establishment of new infections. Thus, its use even after disease is established may prevent or delay further disease development.

Application Rate and Directions: Apply 2.5 lbs product per acre. Dilution rate should be adequate to provide thorough coverage of all leaf and stem tissue. For optimal disease control, arrange nozzles and adjust gallonage to ensure excellent coverage of both sides of leaves and of stems.

For product information call 1-888-6-DUPONT

© 2011 E. I. du Pont de Nemours and Company, 1007 Market Street, Wilmington, Delaware 19898

volume airblast sprayers, and 3-10 gallons per acre for aerial application. Rates of product per acre must be the same for dilute and concentrated sprays.

Add DuPont™ ManKocide® fungicide/bactericide slowly to water in the spray tank with agitation, or premix thoroughly in separate holding tank for concentrate for aircraft sprayers. Continuous agitation is recommended to keep the product in suspension. If needed, adjuvants of the spreader, sticker, or compatibility agent type that are approved for use on growing crops may be used.

PRECAUTIONS AND RESTRICTIONS

- Do not apply more than 16.5 lbs ManKocide® per acre per season.
- If ManKocide® is applied in a spray solution having a pH of less than 6.5, phytotoxicity may occur.
- This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.
- Environmental conditions such as extended periods of wet weather, acid rain, etc. which alter the pH of the leaf surface may affect the performance of DuPont™ ManKocide® fungicide/bactericide resulting in possible phytotoxicity or loss of effectiveness.
- Pesticides may perform in an unpredictable manner when tank mixed especially where several products are involved. Reduced effect on pests or crop injury may occur. Unless recommended by a state/local expert, or the user has small scale direct experience, tank mixing should not be undertaken.
- Do not apply this product via chemigation
- Mixing of this product with products containing diazinon, fosetyl-al or other aluminum containing products, or thio-phanate-methyl is not recommended because of physical incompatibility.
- It must be determined in the selection process if proper application equipment is available and if the waste associated with its use can be properly handled. Materials used in the construction of application equipment is also an important factor as pesticides are often reactive with soft metals such as aluminum and even some synthetic materials such as plastics, rubbers, etc. Therefore it is necessary when working with equipment containing these materials that they are thoroughly flushed with clean water after each days use.
- Carrot seed may not be distributed for human consumption or animal feed. No portion of the carrot seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.

See EPA registered label for additional information.

The DuPont Oval Logo; DuPont™ and ManKocide® are trademarks or registered trademarks of E. I. du Pont de Nemours and Company

Read the Limitation of Warranty and Liability on the product label before buying or using this product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the product label.

R-1117 090811 08-10-11