

FIFRA 24(C)



SPECIAL LOCAL NEED (SLN) LABEL

SePRO Corporation 11550 N. Meridian St., Suite 600, Carmel, IN 46032 USA

(FOR DISTRIBUTION AND USE ONLY IN THE STATE OF IDAHO)

This SLN label for Natrix expires and must not be used or distributed after December 31, 2016

Natrix®

For Use to Control Invasive/Exotic Aquatic Mussels, Snails, Oysters or Clams

Active Ingredient

Copper Ethanolamine Complex† (Mixed CAS#'s 82027-59-6 & 14215-52-2)28.2%

Other Ingredients71.8%

TOTAL100.0%

†Metallic Copper equivalent = 9.1%

EPA Reg. No. 67690-9

EPA SLN No. ID-110007a

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application when applying this product to control invasive/exotic aquatic mussels, snails, oysters or clams.
- Read the label affixed to the container before applying. Carefully follow all precautionary statements and applicable use directions except as specified here.

DIRECTIONS FOR USE

Natrix may be applied to control invasive/exotic aquatic mussel, snail, oyster or clam species, such as: zebra mussels (*Dreissena polymorpha*), quagga mussels (*Dreissena rostriformis bugensis*), Asian clams (*Corbicula fluminea*), and island applesnails (*Pomacea insularum*). Apply at a concentration up to 1.0 ppm of copper in areas containing an active potable water intake; and up to 2.5 ppm of copper in non-potable water or in areas containing potable water intakes that can be turned off until copper concentrations are less than 1.0 ppm at the intake.

Rates and exposure times will vary based on the species and life stage of the species being controlled, as well as treatment conditions. Use the lowest rate and frequency of application necessary to control the target invasive/exotic species. For effective control of adult life stages and depending on site conditions, repeat applications may be necessary. For best results in flowing water, maintain an injection rate for a minimum of 24 hours. Do not treat more than one-half of the lake or pond at one time to avoid depletion of oxygen levels.

- In potable water with an active water intake in the treated area, apply Natrix at a rate up to 3 gallons per acre foot in static water or at a rate up to 32 fluid ounces per cubic feet per second (cfs) per hour in flowing water.

©Copyright 2014 SePRO Corporation

®Natrix is a registered trademark of SePRO Corporation

FPL20140424