RESTRICTED USE PESTICIDE

DUE TO POTENTIAL SECONDARY TOXICITY TO NONTARGET ORGANISMS

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

rozol® VOLE BAIT

FOR CONTROL OF VOLES AND DEER MICE IN HOP FIELDS (POST-HARVEST OR DORMANT-SEASON USE ONLY).

EPA SLN No. ID-1800XX **EPA Reg. No. 7173-242**

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2019.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions and precautions on the EPA registered label.

DIRECTIONS FOR USE:
This product may only be used to control voles (*Microtus* spp.) and deer mice (*Peromyscus* spp.) in hop fields (post-harvest or dormant-season use only).

APPLICATION DIRECTIONS:
Before application, locate vole trails, runway systems and harborage areas to be treated.

Hand spot baiting: Place 1-1/2 ounces (6 tablespoons) of bait in each active hole, trail or runway. Cover each placement with grass or shingle to avoid exposing nontarget organisms, or place bait in a tamper resistant bait station.

Ground or Aerial broadcast baiting: Broadcast evenly at 10 pounds of bait per acre using commercial bait dispensing equipment. Concentrate baiting in areas with heaviest vegetative cover. Do not broadcast bait on bare ground. Do not apply bait in piles or permit piles to be formed by equipment. If populations are high, make a second application 1 to 2 months after the first.

RESTRICTIONS/PRECAUTIONS:
- Do not apply more than 10 pounds of bait per acre per application.
- Do not apply more than 20 pounds of bait per acre per season.
- Do not apply bait during the growing season.
- Only apply bait after fall harvest, before spring growth and when no rain or snow is expected for three consecutive days.
- Do not apply within 10 feet of any body of surface water or where raptors are actively feeding on voles.
- Keep away from humans, domestic animals and pets during application, handling and storage.
- Do not apply bait in a manner that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Keep all other persons out of the treated area during application.
- This product is extremely toxic to mammals and birds. This SLN label should not be used where impact on listed threatened or endangered species is likely. Consult the Federal label for additional restrictions and precautions to protect fish and wildlife.

24(c) registrant:
Liphatech, Inc.
3600 W. Elm Street
Milwaukee, WI 53209
(800) 351-1476

LIPHATECH®

08/15/2018