

# RESTRICTED USE PESTICIDE

## DUE TO POTENTIAL SECONDARY TOXICITY TO NONTARGET ORGANISMS

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

## Section 24(c) Special Local Need Label

FOR USE ONLY WITHIN BEAR LAKE, BINGHAM, BUTTE, CLARK, CUSTER, FREMONT, JEFFERSON, LEMHI, AND TETON COUNTIES IN THE STATE OF IDAHO

# ROZOL<sup>®</sup> GROUND SQUIRREL BAIT

**FOR CONTROL OF WYOMING GROUND SQUIRRELS (*Urocitellus elegans*) IN VINEYARDS, ORCHARDS AND GROVES, TREE AND FORESTRY PLANTATIONS, NURSERIES, RANGELAND, FALLOW AGRICULTURAL LAND, AND NONCROP AREAS INCLUDING BORDER AREAS, BUFFER STRIPS AND CENTER PIVOT CORNERS ADJACENT TO CROPS**

EPA SLN No. ID-190009

EPA Reg. No. 7173-305

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2024.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions and precautions on the EPA registered label.

### DIRECTIONS FOR USE:

This product may be used only for control of Wyoming ground squirrels (*Urocitellus elegans*) in vineyards, orchards and groves, tree and forestry plantations, nurseries, rangeland, fallow agricultural land, and noncrop areas including border areas, buffer strips and center pivot corners adjacent to crops. This product must not be applied directly to food or feed crops except as specified on this label.

**APPLICATION DIRECTIONS:** Ground squirrels prefer to eat fresh green grass and other plants. When these are not available (early spring or when green vegetation dries up in the hot summer), the squirrel's diet converts to grain-based foods and they will readily consume this grain-based bait. Apply bait by broadcast or spot baiting only when fresh green ground cover is not present. Monitor the area to ensure that nontarget species are not consuming the bait or squirrels.

**Bait station baiting:** Apply bait in secured tamper-resistant bait stations. Fill bait stations with 1 to 4 pounds of bait, and secure the stations near active burrows, runways, and other locations where signs of ground squirrel activity are observed. Space stations 20 to 100 feet apart, depending on the density of the squirrel population. Check stations at least weekly to ensure an uninterrupted supply of fresh bait for at least 30 days or until all signs of squirrel feeding have stopped. When feeding by squirrels ceases, and there is no longer a threat of continued reinvasion, remove bait from bait stations and dispose of properly. Bait stations may be kept in place permanently, but should be loaded with toxic bait only when ground squirrels are readily accepting grain-based foods. Do not exceed 4 pounds of bait per station.

**Spot baiting:** Scatter ¼ cup of bait (approximately 2 ounces) over 40 to 50 square feet (a circle approximately 7 to 8 feet diameter) near burrows and active runways. Do not pile the bait. Make a second application 3 to 4 days after the first, following the same procedure as the first application. If squirrel activity persists, a third application may be made 3 to 4 days later. An uninterrupted supply of bait should be available for 6 to 8 days. Do not exceed 10 pounds of bait per acre per application. Do not exceed 20 pounds of bait per acre per year.

**Broadcast baiting:** Using a commercial spreader, apply up to 10 lbs of bait per swath acre, in the area of squirrel activity. Make a second application 4 days after the first. If necessary, make a third application four days after the second application, in order to maintain an uninterrupted supply of bait for 6 to 8 days, but do not exceed 10 pounds of bait per acre per application. Do not exceed 20 pounds of bait per acre per year.

**Surveillance and Follow-Up:** Collect and dispose of dead ground squirrels and search for non-target animals

4 days after the first application and at subsequent intervals of 1 to 2 days for at least 2 weeks after the last bait application, or longer if carcasses are still being found. Wearing gloves, collect and properly dispose of visible carcasses of ground squirrels and other species found in or near treated areas. Dead animals buried on site must be buried a minimum of 18 inches below the ground surface, preferably deeper. Wear gloves and use leakproof plastic bags or other suitable containers for transporting carcasses not buried on site.

**RESTRICTIONS/PRECAUTIONS:**

- Do not apply this product if migratory birds are visiting the application area.
- Do not apply this product by any method not specified on this label.
- Do not apply this product directly to water or within 10 feet of surface water (i.e. streams, rivers, lakes).
- Keep away from humans, domestic animals and pets during application, handling and storage.
- This product is extremely toxic to mammals and birds. This SLN label should not be used where impact on listed threatened or endangered species is likely. Consult the Federal label for additional restrictions and precautions to protect fish and wildlife.

24(c) registrant:  
Liphatech, Inc.  
3600 W. Elm Street  
Milwaukee, WI 53209  
(800) 351-1476

**LIPHATECH®**

05/29/2019