

On-Farm Composting of Dairy Cattle Mortalities



Why Consider Composting?

As knackery and rendering services have declined in many parts of Victoria over the last five or so years, the question of what to do with dead stock has become increasingly problematic for some dairy farmers. There are stringent guidelines for the burial of dead stock (see EPA Victoria Publication 660, www.epa.vic.gov.au). Unfortunately, a few desperate farmers have been known to leave their animals in the 'back paddock' or even dump them in waterways (Fig 1). Dumping of farm animals in this manner is a biosecurity and environmental hazard – it is also illegal and EPA Victoria has prosecuted those responsible for it in the past.

Composting of dead farm animals is a relatively simple and effective process for the routine disposal of dead stock of all sizes (i.e. from poultry to mature cattle). Some farmers have been composting for a long time. The resulting compost can be applied to cultivated land, as top dressing on pastures, or on non-grazed farm areas such as shelterbelts. Compost is a soil conditioner that is able to improve a range of different soil properties, e.g. organic matter content, and provide slow release nutrients.



Figure 1: Illegal dumping of stock is a biosecurity and environmental hazard

What is mortality composting?

Composting is a biological process in which naturally occurring microorganisms and soil animals convert organic materials into a soil-like material called compost. More specifically, mortality composting can be described as the above-ground burial of dead stock in a mound of sawdust, shavings or similar supplemental carbon material. Sufficient supplemental carbon is required around the carcass to absorb bodily fluids and to prevent odours from escaping from the pile.

Mortality composting is generally conducted in 3 stages. In the first stage of composting, the pile is left undisturbed as soft tissue decomposes and bones partially soften. The compost is usually then moved, turned or mixed to begin the second stage, during which time the remaining materials (mainly bones) break down further. Following completion of the second stage, the composting process is completed during a curing or storage phase.

The first (and sometimes second) stage of composting is characterised by high temperatures (>55°C); these conditions result in the elimination of nuisance odours and destruction of pathogens (disease-causing organisms) and weed seeds.

Warning!

If there is suspicion that the animal may have died from a notifiable or exotic disease, the carcass must not be composted but disposed of under the direction of the Victorian Department of Primary Industries. If in doubt, a vet should be contacted.

Selecting the Site for Composting

The site should be located in an elevated area with soils that have low permeability. It may be possible to use an existing hardstand area or a disused concrete pad for composting dead farm animals. If neither is available, a hardstand area that provides all-weather access may need to be established. Instructions for constructing a hardstand are available on the Internet (e.g. www.epa.sa.gov.au/pdfs/cattle.pdf).

Additional requirements include the following:

- The watertable at the site must be at least 1–2 m below the surface;
- The site must not be within 100 m of surface waters (e.g. streams, lakes, wells etc);
- The site should have an adequate slope (1–3%) to allow proper drainage and prevent pooling of water following a rain event;
- Run-off from the composting facility (e.g. during a heavy rainfall event) should be directed to an effluent pond or a vegetative filter strip / infiltration area;
- The site should have all-weather access and have minimum interference from other traffic;
- Consideration should be given to prevailing winds and the location and proximity of the site in relation to the farm house and neighbours in order to prevent potential odour problems;
- Fence off the site to help eliminate scavenging animals and other livestock accessing the piles.



Figure 2: Establish 45-60 cm base layer of carbon material



Figure 3: Place the carcass in the centre of the base



Figure 4: Cover the carcass with 60 cm carbon material

How is Composting Done?

These instructions are for composting dairy cattle using the open pile method, which is likely to be the method of choice for most farmers.

Constructing the Pile

The construction of the pile requires considerable quantities of carbon material; about 10 - 12m³ is needed to establish a composting pile for a 450-kg carcass. Up to 50% of carbon material can be comprised of finished compost from previous composting piles.

Follow this process for constructing a mortality composting pile:

Step 1. Establish a base layer of relatively dry carbon material at the composting hardstand area (Fig. 2). The depth of the base layer should range between approximately 45 cm for small carcasses and 60 cm for large carcasses. The dimensions of this base layer must be large enough to accommodate the carcasses with >60 cm space around the edges.

Step 2. Place the carcass in the centre of the base layer (Fig. 3).

Step 3. Cover the carcass with carbon material to a depth of about 60 cm (Fig. 4). The material used to cover the animal should be damp to enhance the composting process. The material must feel moist but not be too wet. Composting material is too wet when water can be squeezed from it (droplets appear between fingers).

Step 4. Leave the pile undisturbed for at least 4 months. Provided that the animal is covered well enough, scavengers should not be attracted to the pile.

Step 5. When the soft tissue has completely decomposed after 4-6 months the pile can be turned and watered (Fig. 5) to complete the composting process.

Step 6. When the pile starts to cool down, leave it to cure for at least another 4 weeks before using the compost. It is not necessary to turn the pile during the curing stage.

Although the composting process is complete, large bones of adult cattle such as the skull, the spine, shoulder blades or leg bones may still be present. However, they are usually 'clean' (free of soft tissue) and have become brittle, starting to disintegrate (Fig 6).



Figure 5: Watering the compost at turning

Monitoring the Composting Process

1. Monitor compost pile temperatures (> 60 cm depth) weekly for the first four weeks after establishing the pile and after turning (Fig. 7). Subsequently, fortnightly measurements should be sufficient. Take temperature readings (three are suggested) from several points in the pile and record them down for future reference.
2. After establishing the pile, visit the pile daily for the first week, to ensure there is no odour or liquid running from the base of the pile, and no dogs or foxes have dug into the pile to get access to the carcass. Add more carbon material if needed to cover the animal or to mop up seeping fluids.
3. From about 4 months (less for younger, smaller animals), periodically check for signs of remaining soft tissue. If none remains, the pile can be turned and watered as described above.



Figure 6: Bones remaining after 4 months composting

Biosecurity

Composting is known to control nearly all pathogens - viruses, bacteria, fungi, protozoa (including cysts) and helminth ova to acceptably low levels. Exceptions to this are some spore-forming bacteria (e.g. *Bacillus anthracis*, 'anthrax') and prions like BSE (bovine spongiform encephalopathy). Prions are highly resistant to both physical and chemical means of inactivating pathogens and for this reason it is assumed that composting will be ineffective in reducing infectivity of prion-infected carcasses.

Pathogens are killed during composting by multiple means, such as high temperatures, the direct and indirect effects of other microorganisms and the presence of organic acids and ammonia in the compost. Not only is temperature considered to be the most important factor in killing pathogens, it is also relatively easy to measure during composting.

The heat required for the inactivation of pathogens is a function of both temperature and length (time) of exposure. Exposure to an average temperature during composting of 55 to 60°C for a couple of days is usually sufficient to kill the vast majority of pathogens.



Figure 7: Monitoring temperature during composting




Figure 8: Temperature variation within a typical pile

To achieve efficient pathogen kill, all materials in a compost pile must be exposed to high temperatures for prolonged periods. In piles, there is great variation in the temperature profile from the cool outside layers to the hot central mass (Fig. 8). As a result, piles are usually turned periodically to expose the outer layers of the pile to high temperature composting. In mortality composting, piles are turned after the soft tissue has decomposed (stage 2).

Risk Management

Proficient design and operation of mortality composting systems is very important to ensure that all material achieves adequate temperatures for long enough to kill pathogens. This is achieved for mortality composting piles provided that temperature is monitored.

The current state of knowledge suggests that taking the following factors into consideration will reduce the potential biosecurity risk associated with mortality composting:

- Attention to site design and layout to minimise scavenging and contamination of ground and surface water with pathogens;
- Using a minimum two-stage composting system followed by the use of a curing phase to properly complete the composting process;
- Where possible, fully encapsulating mortalities in 'clean' carbon source; use sufficient carbon source to absorb liquids and odorous gasses produced during composting;
- Monitor and manage the composting process to maximise progress towards the full completion of composting (e.g. temperature, monitoring, watering and turning);
- Attention to basic standards of hygiene (e.g. minimising pooling of water at the site, regular sanitising of equipment and keeping it separate from production facilities, use of personal safety equipment by compost operators).

Health and Safety

A compost pile contains living microorganisms including moulds, bacteria, fungi and protozoa. These microorganisms can cause adverse reactions, particularly in people with a weakened immune system, asthma or a punctured ear-drum, or if somebody takes antibiotics or adrenal cortical hormones. Dust particles or bioaerosols released from the pile during handling of the compost on rare occasions may cause skin irritations, eye infection or respiratory illness.

The following precautions should always be followed when handling dead stock or compost materials:

- Wear gloves;
- Wash hands after handling dead stock or compost;
- Avoid breathing dusts or mists;
- Wear particulate mask;
- Wear dust resistant eye protection;
- Wash work clothing regularly;
- Prevent injury when operating machinery.

Further information

Dr Kevin Wilkinson, Department of Primary Industries, PO Box 4166, Parkville, Victoria, 3052. Ph. 03 8341 2412.

Email: kevin.wilkinson@dpi.vic.gov.au

Johannes Biala, The Organic Force, PO Box 74, Wynnum, Qld, 4178. Ph. 07 3901 1152. Email: biala@optusnet.com.au

Detailed instructions for mortality composting available on the Dairying for Tomorrow website (www.dairyingfortomorrow.com)

Published by: Department of Primary Industries
Future Farming Systems Division
PO Box 4166, Parkville, Victoria, 3052.
September, 2007

© The State of Victoria, 2007

This publication is copyright. No part may be reproduced by any process except in accordance with the provisions of the Copyright Act 1968.

Authorised by: Victorian Government
1 Spring St
Melbourne, Victoria 3000, Australia

Disclaimer

This publication may be of assistance to you but neither the State of Victoria and its officers guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purposes and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on any information in this publication. 4

For more information about DPI visit the website at www.dpi.vic.gov.au or call the Customer Service Centre on 136 186.