

RESTRICTED USE PESTICIDE

DUE TO HAZARDS TO NONTARGET SPECIES

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

FIFRA SECTION 24(c) SPECIAL LOCAL NEED LABEL

FOR DISTRIBUTION AND USE ONLY WITHIN BUTTE COUNTY, IDAHO.

This product may be used to control Columbian ground squirrels and Wyoming ground squirrels in alfalfa and barley hay mix.

This label is valid until December 31, 2029, or until otherwise amended, withdrawn, canceled, or suspended.

ZINC PHOSPHIDE ON OATS

Alternate Brand Name: ZINC PHOSPHIDE ON OATS – ID

EPA Reg. No. 56228-14

EPA SLN No. ID-190001

ACTIVE INGREDIENT

Zinc Phosphide..... 2.00%

OTHER INGREDIENTS 98.00%

TOTAL..... 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Misuse may result in civil or criminal enforcement action.

READ THIS LABEL: Read this entire label and the label for EPA Registration Number 56228-14. This product must be used strictly in accordance with this label's precautionary statements and use directions on both labels, as well as with all applicable State and Federal laws and regulations.

This state-specific Section 24(c) labeling and the Section 3 labeling for EPA Reg. No. 56228-14 must be in possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on this label and the EPA-registered label for EPA Reg. No. 56228-14.

Before using this product, contact the U.S. Fish and Wildlife Service and the applicable State wildlife agency and obtain all necessary kill or collecting permits. Use only for the sites, pests, and application methods described on this label.

APPLICATION DIRECTIONS

For control of Columbian ground squirrels (*Urocyonotellus columbianus*) and Wyoming ground squirrels (*Urocyonotellus elegans*).

PREBAITING: Prebaiting with nontoxic oats is required to improve acceptance of this zinc phosphide bait. Prebait should be applied, at rates appropriate for toxic bait, 2 or 3 days before toxic baits are used.

Application directions and site-specific use restrictions for this label are contained in Table 1.

Table 1.

Use Site	Use Restrictions	Application Directions
Alfalfa	All applications must occur shortly after a cutting of the hay, and/or prior to the next growth's attaining a length of 2 inches. Alfalfa forage from treated areas must not be harvested until it reaches maturity.	BAITING: Broadcast Baiting: This product may be broadcast by air or ground-driven dispensing devices. Apply toxic bait at a rate up to 10 lbs/A (0.18 lb ai/A). A second application may be made after 10 days. Do not apply in piles or permit piles to be formed by equipment. Apply toxic bait at a maximum rate of 20 lbs/A (0.4 lb ai/A) per year.
Barley Hay Mix	Do not apply this product to barley within 50 days of harvest. Do not graze animals in treated areas.	BAITING: Broadcast Baiting: This product may be broadcast by air or ground-driven dispensing devices. Apply toxic bait at a rate up to 6 lbs/A (0.12 lb ai/A). A second application may be made after 25 days. Apply toxic bait at a maximum rate of 12 lbs/A (0.24 lb ai/A) per year. All applications must be made prior to the boot stage.

FOLLOW-UP: Wearing waterproof gloves, collect and dispose of dead, exposed animals. Dispose of leftover toxic bait in accordance with **PESTICIDE DISPOSAL** instructions under **STORAGE AND DISPOSAL**. If reinfestation does occur, repeat treatment.

24(c) Registrant:

U.S. Department of Agriculture
Animal and Plant Health Inspection Service
4700 River Road, Unit 149
Riverdale, MD 20737

Local Contact:

USDA APHIS Wildlife Services
9134 West Blackeagle Drive
Boise, ID 83709

Net Contents: _____

Batch Code: _____