

# FIFRA Section 24(c) Special Local Need (SLN) Label



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Stinger<sup>®</sup>

EPA Reg. No. 62719-73

**Postemergence Control of Certain Noxious and Broadleaf Weeds in Hops**  
(For Distribution and Use Only in the State of Idaho)

24(c) Special Local Need Registration EPA SLN No. ID-230003

This label is valid until December 31, 2029, or until otherwise amended, withdrawn,  
cancelled, or suspended.

### Manufacturer's Conditions and Risks of Use for Special Local Need

USE OF Stinger<sup>®</sup> herbicide (THE "PRODUCT") ON HOPS (THE "CROP") IN ACCORDANCE WITH THIS SPECIAL LOCAL NEED LABELING MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED BEFORE USING THE PRODUCT ON THE CROP.

Corteva Agriscience makes the Product available for use in the manner described in this Special Local Need Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury, Crop yield reduction or Crop loss. The decision to use this Product in the manner described in this Special Local Need Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this Special Local Need Labeling, you acknowledge and accept that:

- (1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- (2) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Special Local Need Labeling;
- (3) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaims all warranties, express or implied, including any implied warranty of merchantability;
- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;

(5) these Conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Corteva Agriscience, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

**If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.**

**These Conditions and Risks of Use for Special Local Need are required by Corteva Agriscience and not specified by the U.S. EPA or the State of Idaho.**

### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This Special Local Need Labeling must be in the possession of the user at the time of application.
- Read the label affixed to the Product container before applying Product on the Crop. Carefully follow all use precautions and limitations.
- In addition to restrictions and limitations imposed by this Special Local Need Labeling, use of Product on the Crop is also subject to all use precautions and limitations imposed by the label affixed to the Product container.

### Directions for Use

Refer to the product label affixed to the container for General Use Precautions, Mixing and Application instructions, and a complete list of weeds controlled or suppressed.

#### Target Broadleaf Weeds, Application Rate, and Use Restrictions:

Target Broadleaf Weeds	Application Rate (pint/acre)	Use Restrictions
dock, curly groundsel, common horseweed knapweeds <sup>†</sup> lettuce, prickly sowthistles <sup>†</sup> starthistle, yellow thistle, Canada	1/3 – 2/3 (0.125 – 0.25 lb ai/acre)	<ul style="list-style-type: none"><li>• Apply after training when the growing points of the hop plants are above the spray area.</li><li>• <b>Do not</b> make more than 2 broadcast applications per crop per year.<sup>††</sup></li><li>• <b>Preharvest Interval:</b> Do not apply within 30 days of harvest.</li><li>• <b>Re-treatment Interval:</b> 21 days</li></ul>

<sup>†</sup>For perennial weeds such as Russian knapweed and perennial sowthistle, Stinger will control the top growth and inhibit regrowth during the season of application (season-long control). At higher use rates shown on this label, Stinger may cause a reduction in shoot regrowth in the season following application; however, plant response may be inconsistent due to inherent variability in shoot regrowth from perennial root systems.

<sup>††</sup>The total combined usage of Stinger from all types of applications must not exceed a total of 2/3 pint (0.25 lb ai clopyralid) per crop per year.

**Advisory:** Some transient minor leaf cupping may occur to lower leaves and suckers if spray comes into contact with plant. Different varieties of hops may exhibit differing levels of sensitivity to herbicides. It is recommended that applicators test a small area before applying to entire hop yard. Extreme growing conditions such as drought prior to, at, or following time of application may increase the risk of crop injury at all stages of growth.

**Broadcast Application Rates:** Apply uniformly with ground equipment in 10 to 20 gallons of water per acre. A second application may be made as long as the total amount applied does not exceed 2/3 pint per acre of Stinger per crop per year. For control of Canada thistle, apply Stinger after the majority of basal leaves have emerged but prior to bud stage.

**Hand-Held Sprayers:** Hand-held sprayers may be used for spot applications of Stinger if care is taken to apply the spray uniformly and at a rate equivalent to a broadcast application (example: 0.25 fl oz or 7.3 ml of Stinger per gallon of water is equal to the broadcast rate of 2/3 pint of Stinger per acre). To prevent misapplication, spot treatments should be applied with a calibrated boom or with hand sprayers according to the directions in the Hand-Held Sprayers section of the instructions for spot application on the container label for Stinger.

**Tank Mixtures:** Stinger may be tank mixed with other herbicides that are registered and labeled for use on hops. Follow the Directions for Use of the labeling for any tank mix partner used in tank mixture with this product.

**Specific Use Restrictions:**

- **Chemigation:** Do not apply Stinger through any type of irrigation system.
- **Do not** apply by aircraft.
- **Do not** apply within 6 to 8 hours of expected rainfall or irrigation.
- **Do not** compost treated vegetation if compost will be used on sensitive plants.

<sup>TM®</sup> Trademarks of Corteva Agriscience and its affiliated companies

R043-218

Accepted: \_\_/\_\_/\_\_

Replaces R043-172