



Section 24(c) Special Local Need Label

S-METOLACHLOR

GROUP

15

HERBICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

Dual Magnum® Herbicide
Dry Bulb Onion

Post-Emergent Application for Residual Weed Control
Fall Preplant Application for Pre-emergent Control or Suppression of Yellow Nutsedge

EPA Reg. No. 100-816
EPA SLN No. ID-990016

This label expires and must not be distributed or used in accordance with this SLN registration
after December 31, 2029

SYNGENTA'S SPECIAL CONDITIONS, RISKS OF USE AND DISCLAIMER FOR USE OF DUAL MAGNUM HERBICIDE ON CROPS ON THIS 24(c) LABEL

IMPORTANT- READ BEFORE USE

THESE CONDITIONS RISKS OF USE AND DISCLAIMER ARE REQUIRED BY SYNGENTA CROP
PROTECTION LLC AND NOT SPECIFIED BY U.S. EPA OR THE OF STATE OF IDAHO

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SYNGENTA CROP PROTECTION, LLC INTENDS THAT THE PRODUCT THAT IS THE SUBJECT OF THIS SECTION 24(c) LABEL BE PURCHASED ONLY BY END USERS WHO AGREE BY ELECTRONIC SIGNATURE ON SYNGENTA CROP PROTECTION'S INTERNET SITE TO THE TERMS AND CONDITIONS REQUIRED BY SYNGENTA CROP PROTECTION, LLC INCLUDING A WAIVER AND RELEASE FROM ALL LIABILITY AND INDEMNIFICATION BY THE USER AND/OR GROWER OF SYNGENTA AND OTHERS FOR FAILURE TO PERFORM AND FOR CROP INJURY, CROP YIELD REDUCTION, AND/OR CROP LOSS FROM USE OF DUAL MAGNUM HERBICIDE ON CROPS ON THIS 24(c) LABEL. IF SUCH TERMS AND CONDITIONS ARE UNACCEPTABLE, RETURN THE DUAL MAGNUM HERBICIDE AT ONCE UNOPENED OR USE THE DUAL MAGNUM HERBICIDE FOR A DIFFERENT APPROVED USE IN ACCORDANCE WITH THE LABEL AFFIXED TO THE PRODUCT CONTAINER.

USE OF DUAL MAGNUM HERBICIDE (THE "PRODUCT") ON CROPS LISTED (THE "CROP") FOR THIS SPECIAL LOCAL NEED MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS AS FURTHER DISCUSSED BELOW. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED BEFORE USING THE PRODUCT ON THE CROP. SYNGENTA RECOMMENDS THAT THE USER TEST THIS PRODUCT TO DETERMINE ITS SUITABILITY FOR SUCH INTENDED USE.

Syngenta Crop Protection, LLC makes the Product available for use in the manner described in this 24(c) Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury, Crop yield reduction or Crop loss.

The decision to use this Product in the manner described in this 24(c) Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors. Syngenta recommends that the user test this Product to determine its suitability for such intended use.

By purchasing the Product for use, or using the Product in the manner described in this 24(c) Labeling, you acknowledge and accept that, to the extent consistent with applicable law:

- 1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- 2) Syngenta Crop Protection, LLC do not make, and do not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this 24(c) labeling;
- 3) Syngenta Crop Protection, LLC do not make, and do not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of merchantability;
- 4) Syngenta Crop Protection, LLC disclaim all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- 5) these conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Syngenta Crop Protection, LLC or their respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.
- The risk of crop injury increases when adjuvants (e.g., non-ionic surfactants, crop oils, etc.), nitrogen sources (e.g., AMS, UAN), fertilizers or other pesticides are applied with Dual Magnum Herbicide.

Dual Magnum Herbicide will not control emerged weeds. For effective weed control, Dual Magnum Herbicide must be applied to clean-tilled soil or to ground where emerged weeds have been controlled by another ns.

Do not apply under conditions which favor runoff or wind erosion of soil containing this product to nontarget areas.

To prevent off-site movement due to runoff or wind erosion:

1. Avoid treating powdery dry or light sand soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.
2. Do not apply to impervious substrates, such as paved or highly compacted surfaces.
3. Do not use tailwater from the first flood or furrow irrigation of treated fields to treat nontarget crops, unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.

Dry Bulb Onions – Postemergent Application to the Crop

For suppression and control of yellow nutsedge, grass, and some broadleaf weeds (see **Weeds Controlled** on the Dual Magnum Herbicide label), apply Dual Magnum Herbicide at the two (2) true leaf stage of onions at rates of 0.67 - 1.33 pints (0.64-1.27 lb active ingredient) per acre, depending on soil type and target weed. Use the lower rate on light, sandy soils and where a general weed spectrum is targeted. The higher rate will provide improved yellow nutsedge control, but comes with an increase risk of crop injury. One additional application of 0.67-1.33 pints may be applied 21 days or more after the first treatment, if needed, provided no fall preplant applications of Dual Magnum Herbicide were made. Dual Magnum Herbicide provides good to excellent control of yellow nutsedge. **If nutsedge is not a target weed delaying Dual Magnum Herbicide application until onions have three true leaves may reduce the risk of crop injury.**

Field research in some western states has shown that adverse weather conditions such as excessive rain or cold, wet conditions after application can increase the risk of crop injury from Dual Magnum Herbicide. Stand thinning, stunting and yield loss may occur.

Onion tolerance to Dual Magnum Herbicide increases with increasing onion size. However, growers must weigh the need to control early nutsedge flushes with the potential risk of crop injury.

Restrictions (1) **Do not** apply within 60 days of harvest. (2) **Do not** harvest green onions. (3) **Do not** apply this product through any types of irrigation system. (4) **Do not** graze animals on green forage or stubble. (5) If a fall preplant application of Dual Magnum Herbicide was used for nutsedge, only one post-emergent application at a maximum rate of 1.33 pints/A is allowed. (6) **Do not** apply more than 2.68 pints per acre (2.55 lb ai/A) to dry bulb onions as a combined total across all application timings and use patterns to produce that crop. Do not apply more than 2.55 lb ai/A per year of S-metolachlor-containing products.

Dry Bulb Onions – Fall Preplant Application

For pre-emergent control or suppression of yellow nutsedge the following spring in dry bulb onions apply 1.33 pt/A of Dual Magnum Herbicide in the fall after the harvest of the previous crop but before freeze-up. Fall applications of Dual Magnum Herbicide can be surface-applied or incorporated. To reduce the risk of crop injury, apply at least 100 days prior to the planting of onion (seed, sets, or transplants).

Precautions: 1) In general, the risk of crop injury is greater on lighter textured soils and with higher application rates. 2) The addition of another registered herbicide as a tank mixture or in a program with Dual Magnum Herbicide can increase the risk of crop injury.

Restrictions: (1) Make no more than one fall application per crop. (2) Apply not more than 1.33 pt/A in a single fall preplant application. (3) Do not apply this product, for this use, through any types of irrigation system. (4) Do not apply to frozen ground.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

Ground Water Advisory

S-metolachlor is known to leach through soil into ground water under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water or through ground spray drift. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several weeks or months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of S-metolachlor from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Non-Target Organism Advisory:

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Care must be taken when using this product to prevent back siphoning into wells, spills, or improper disposal of excess pesticide, spray mixtures, or rinsates. Check-valves or antisiphoning devices must be used on all mixing and/or irrigation equipment.

©2024 Syngenta

Dual Magnum® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, NC 27419-8300

Label Code: ID0816047FA1024