FIFRA 24(c) Special Local Need Label (SLN)
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

ISOCYCLOSERAM	GROUP	30	INSECTICIDE

Vertento®
For Control of Plant Bug on Alfalfa Grown for Seed

EPA Reg. No. 100-1711
EPA SLN No. ID-260007

Active Ingredient:	
Isocycloseram**	34.8%
Other Ingredients:	65.2%
Total:	**100.0%**

**PLINAZOLIN® technology denotes the Syngenta trademark for the active ingredient isocycloseram*
**CAS No. 2061933-85-3

A21550 CP is formulated as a suspension concentrate and contains 3.33 lb of isocycloseram per gallon.

KEEP OUT OF REACH OF CHILDREN
CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and main EPA-registered label for Vertento, EPA Reg. No. 100-1171 must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the main EPA-registered label for Vertento, EPA Reg. No. 100-1171.

FOLIAR APPLICATION

Alfalfa			
Target Pest	**Rate (fl oz/A)**	**Application Timing**	**Use Directions**
Tarnished plant bug (*Lygus* sp.)	1.6 – 2.0	Time applications to the most susceptible insect pest life-stage at locally determined action thresholds before populations reach damaging levels.	Under high pest populations, apply a higher rate within the labeled rate range. Apply this product diluted in a minimum volume of 10 gal/A by ground. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
Resistance Management: • Refer to **Section 3.2.** • If Vertento is used on any given cutting, do not use Vertento or any other isocycloseram-containing product on the subsequent cutting. Rotate to a different mode of action before using Vertento or any other isocycloseram-containing product on the following cutting.			
Refer to **Section 6.1** of the Section 3 label for additional product use restrictions.			

RESTRICTIONS AND PRECAUTIONS

- Restricted Entry Interval (REI): 12 hours
- **Maximum Single Application Rate:** 2.0 fl oz/A/application
  a. **DO NOT** exceed 0.0520 lb ai/A of isocycloseram-containing products.
- **Minimum Application Interval:** 7 days
- **Maximum Annual Rate:**
  a. 4.0 fl oz/A/cutting
    i. **DO NOT** exceed 0.104 lb ai/A/cutting of isocycloseram-containing products including all application types (seed treatment, soil, and foliar).
  b. 8.0 fl oz/A/year
    i. **DO NOT** exceed 0.208 lb ai/A/year of isocycloseram-containing products including all application types (seed treatment, soil, and foliar).

- **DO NOT** make more than four applications per year.
- **DO NOT** apply by air.
- **DO NOT** apply by chemigation.
- **DO NOT** apply 3 days prior to flowering until the end of the flowering period.
- **Pre-Harvest Interval (PHI):** 7 days

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

SURFACE WATER ADVISORY

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having a medium potential for reaching both surface water and aquatic sediment via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of isocycloseram from runoff water and sediment. Runoff of this product will be reduced by avoiding application when rainfall or irrigation is expected. Do not make applications during rain and avoid making applications when rainfall is expected before the product has sufficient time to dry.

NON-TARGET ORGANISM ADVISORY

This pesticide is toxic to fish and highly toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not apply when weather conditions favor drift from target areas.

POLLINATOR PRECAUTIONS

This product is highly toxic to bees and other pollinating insects exposed to direct treatment, or to residues in/on blooming crops or weeds. Protect pollinating insects by following label directions intended to minimize drift and to reduce risk to these organisms.

The RT_{25} (Residual Time to 25% mortality; the length of time over which field weathered foliar residues remain toxic to honey bees) for this product is ≤ 3 hours.

CROPS GROWN FOR SEED

(1) Alfalfa seed from treated fields may not be used for growing sprouts for humans or animal consumption. (2) **DO NOT** feed or graze alfalfa forage or fodder. **DO NOT** cut alfalfa for hay or forage. (3) Producers of alfalfa seed who use this product, or cause the product to be used on fields they operate, are required to inform, in writing, conditioners receiving seed produced on fields treated with this product. A copy of this labeling is required to be provided to the conditioner by the producer. Processed seed must be labeled “Not for human or animal consumption” at the processing plant. The processor must dispose of all seed screenings in such a way that they cannot be distributed or used for food or feed.

©2026 Syngenta

Vertento® and the Syngenta logo are trademarks of a Syngenta Group Company.

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: ID1711758AC0426