

FIFRA Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

MIDAC[®] 4

To allow in-furrow application in sugar beets.

Active Ingredient:	By Wt
Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine ..	40.0%
Other Ingredients:.....	60.0%
Total:	100.0%

This product contains 4.0 pounds of active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN
CAUTION

EPA Reg. No. 89118-16

EPA SLN No. ID-260004

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label and the federal label for this product must be accessible to the user at the time of the pesticide application.
- Read the label affixed to the container for Midac 4 before applying.
- Follow all applicable directions, restrictions, and precautions on this Supplemental Label and the main EPA-registered label. It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Use of Midac 4 according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Midac 4.
- **DO NOT** apply this product through any type of irrigation system.

Environmental Hazards

This pesticide is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. **DO NOT** apply this product or allow it to drift to blooming crops or weeds if bees are foraging in the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements in this labeling about personal protective equipment, restricted-entry intervals, and notification to works. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: if the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Coveralls.
- Shoes plus socks.
- Gloves made of: barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, polyvinylchloride (PVC) ≥14 mils or viton ≥14 mils.

SPECIFIC DIRECTIONS FOR USE

SUGAR BEETS	
FIELD SOIL APPLICATIONS	
PESTS	USE RATES fl oz product/A (lb a.i./A)
Sugar Beet Root Maggot	6.0 (0.19)
Field Soil Application Methods: • Make an application as an in-furrow dribble or spray directed on or below the seed during planting.	
Field Soil Use Restrictions: • Application Timing: DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging. • Application Restrictions: ○ DO NOT use on crops grown for seed. ○ DO NOT use on fields planted with imidacloprid treated seed. • Annual Maximum: ○ DO NOT exceed 6.0 fl oz of VCP-034 (0.19 lb a.i.) per acre per calendar year. ○ DO NOT exceed 0.19 lb imidacloprid per acre per calendar year from all imidacloprid containing products.	

At Plant In-Furrow Application Rates (fl oz product per 1000 row ft)						
fl oz product per acre	Average Row Spacing (inches)					
	20	22	24	26	28	30
6.0	0.23	0.25	0.28	0.30	0.32	0.34

