

**Special Local Need** 

### FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

## **ASSAIL® 70WP INSECTICIDE**

#### FOR CONTROL OF LYGUS BUG IN ALFALFA GROWN FOR SEED

This label is valid until December 31, 2024 or until otherwise amended, withdrawn, cancelled or suspended.

#### ACTIVE INGREDIENT: acetamiprid

(E)- N <sup>1</sup> -[(6-chloro-3-pyridly)methyl] – N <sup>2</sup> -cyano-N <sup>1</sup> -methyl acetamidine	70.0% by wt.
OTHER INGREDIENTS	<u>30.0%</u> by wt.
TOTAL	.100.0%

EPA Reg. No. 8033-23-70506

EPA SLN No. ID-070011B

#### **ENVIRONMENTAL HAZARDS**

This product is toxic to wildlife. This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are visiting the treated area. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Do not contaminate water used for irrigation or domestic purposes.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This supplemental label and the federally registered label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this supplemental label and the main EPA registered label.

Spray volume: Apply in a minimum finished spray volume of 20 gallons per acre by ground and 10 gallons per acre by air. To achieve optimum pest control, it is important to obtain thorough and uniform spray coverage.

USE	RATE/ACRE	SPECIFIC DIRECTIONS
Lygus bugs	2.3 – 3.5 oz. (0.101 – 0.153 lb ai)	<ul> <li>Begin application when treatment thresholds have been reached. Use the high rate under heavy insect pressure and for longer residual control.</li> <li>Do not make more than 2 applications per season.</li> <li>Do not apply more than once every 7 days.</li> <li>Do not exceed a total of 0.4 lbs. active ingredient (9.0 ozs. product) per acre per season.</li> </ul>

#### **RESTRICTIONS:**

Do not cut current year's treated alfalfa seed crop for hay or forage. Do not graze current year's treated alfalfa seed crop. Do not use treated alfalfa seed for sprouting.



# **Special Local Need**

#### **RESISTANCE MANAGEMENT:**

Acetamiprid is the active ingredient in ASSAIL<sup>®</sup> 70WP Insecticide. It is a member of a class of chemicals known as neonicotinoids. Rotating ASSAIL<sup>®</sup> 70WP Insecticide with pesticides with different modes of action may delay or prevent development of insecticide resistance. The use of ASSAIL<sup>®</sup> 70WP Insecticide should conform to resistance management strategies established for the use area. Consult your agricultural advisor, university contact or extension service for recommended pest management practices for your area. Use recommended integrated pest management practices so that you are not relying solely on chemical control in your crop production.

ASSAIL® is a registered trademark of Nippon Soda Co., Ltd.

4/20/2020