



**FIFRA Section 24 (c)
Special Local Need**

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

ACEPHATE 97UP INSECTICIDE

EPA Reg. No. 70506-8

EPA SLN No. 10-170005

ACTIVE INGREDIENT	BY WT.
Acephate: (O,S-Dimethyl acetylphosphoramidohioate).....	97%*
OTHER INGREDIENTS	<u>3%</u>
TOTAL	100 %

*U.S. Patent Nos. 6,752,943 and 6,875,381

CAUTION

Keep Out of Reach of Children

FOR INSECT CONTROL IN ALFALFA GROWN FOR SEED ONLY

This label is valid until December 31, 2025 or until otherwise amended, withdrawn, canceled or suspended.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This FIFRA Section 24 (c) Special Local Need (SLN) label and the federal label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label. Read the entire Directions for Use before using this product. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical resistant gloves made of any waterproof material and shoes plus socks.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds.

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash-waters.

Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Eliminations System (NPDES) permitted and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

USE RESTRICTIONS

Do not apply this product through any type of irrigation system.

No portion of this alfalfa seed crop may be used or distributed for human food or animal feed purposes.

This product may pose a hazard to endangered aquatic species. Refer to and follow the spray drift precautions on the US EPA registered label for this product.

Rotational crops may be planted no sooner than 30 days after the last application.

SPECIAL CROP USE RESTRICTIONS

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a non-feed/non-food use. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have established pesticide residue tolerances for this crop. Consequently, no portion of this crop may be used or distributed for human consumption or animal feed. This restriction pertains to, but is not limited to: green chop, forage, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage, or seed screenings, and the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that screenings cannot be distributed or used for food or feed purposes. All restrictions listed in the Idaho Rules Governing Pesticide & Chemigation Use & Application, IDAPA 02.03.03.800 must be followed.

Any seed from a field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

"This seed was produced using one or more products for which the United States Environmental Protection Agency not established pesticide residue tolerances. **This seed, in whole, as sprouts, or in any form, may not be used for human consumption or animal feed.** Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration and other regulatory agencies. "

Alfalfa Grown For Seed For Control of Lygus Bug

PESTS CONTROLLED	RATE	APPLICATION INSTRUCTIONS	COMMENTS
Lygus Bug	Use 1 lb (0.97 lb ai) of Acephate 97UP per acre per application. Do not exceed 2 lbs (1.94 lbs ai) of product per acre per year.	Ground - apply in 10 to 60 gallons of water. Air - apply in 5 to 10 gallons of water per acre. Make one repeat application at no less than 7 day spray interval, if necessary to maintain control.	Applications may be made up to 3 days prior to placing bees in or around the field. Do not make more than 2 applications per acre per year. The REI is 24 hours. Some data indicate that Leafcutter bees and Alkali bees may be more sensitive to acephate exposure than other tested species.

Rev 7-14-2021