



# FIFRA Section 24(c) Special Local Need (SLN)

## FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

For Control of Aphids, Leafhoppers, Tarnished Plant Bug & Western Plant Bug in  
Clover Grown For Seed

### Sivanto™ prime

EPA Reg. No. 264-1141

SLN. No. ID-150002

THIS LABEL IS VALID UNTIL DECEMBER 31, 2025 UNLESS OTHERWISE AMENDED, WITHDRAWN, CANCELED,  
OR SUSPENDED.

**ACTIVE INGREDIENT:**

Flupyradifurone\* ..... 17.09%

**OTHER INGREDIENTS:** ..... 82.91%

**TOTAL:** ..... 100.00%

---

Contains 1.67 pounds active ingredient per U.S. gallon (200 grams AI/liter)

---

## KEEP OUT OF REACH OF CHILDREN CAUTION

---

### ENVIRONMENTAL HAZARDS

---

#### Terrestrial Use

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

#### Non-Target Organisms

This pesticide is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Toxic to adult bees in laboratory studies via oral exposure, however, not toxic to bees through contact exposure, and field studies conducted with this product have shown no effects on honeybee colony development.

#### Ground Water Advisory

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

#### Surface Water Label Advisories

Flupyradifurone and its degradate difluoroacetic acid may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. Flupyradifurone and its degradate difluoroacetic acid are classified as having medium and high potential, respectively, for reaching surface water via run-off for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of flupyradifurone and its degradate difluoroacetic acid from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label and the EPA-registered label for this product must be in the possession of the user at the time of the pesticide application. Follow all applicable directions, restrictions and precautions on this SLN label and the EPA-registered label.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval **(REI) of 4 hours**, with the exception of California and New York. In the state of California the REI is 12 hours. In the state of New York the REI is 4 hours for all listed crops except for grapes for which the REI is 12 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area without restrictions if there will be no contact with anything that has been treated.

**PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:**

- Coveralls
- Chemical resistant gloves
- Chemical resistant footwear plus socks

## CLOVER GROWN FOR SEED

Pests Controlled	Product Rate (fl oz/A)
Aphids Leafhoppers	7.0 – 10.5
Threecornered alfalfa hopper Whiteflies	10.5 – 14.0
Pests Suppressed	
Tarnished plant bug ( <i>Lygus lineolaris</i> ) Western plant bug ( <i>Lygus hesperus</i> )	14.0
<b>Foliar Application Restrictions:</b> Pre-Harvest Interval (PHI): <b>14 days</b> – Clover Minimum interval between applications: <b>10 days</b> Minimum application volumes: <b>10 gallons/Acre</b> (Ground); <b>2 gallons/Acre</b> (Aerial) Maximum SIVANTO prime allowed per year: <b>28.0 fluid ounces/Acre</b> (0.365 lb AI/Acre).	

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.*

*For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

**Bayer CropScience LP**  
800 N.Lindbergh Blvd.  
St. Louis, MO 63167