



**FIFRA 24(c) Special  
Local Need Label**

EPA SLN No.: ID-150008  
EPA Reg. No. 279-3337

**FOR DISTRIBUTION AND USE  
ONLY IN IDAHO**

**This label for SPARTAN CHARGE herbicide is valid until December 31, 2025 or until withdrawn, canceled or suspended.**

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.**

**THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**

**SPARTAN CHARGE Herbicide for Use in Spring Wheat**

**Directions For Use**

**Apply SPARTAN CHARGE herbicide as a broadcast treatment, post plant pre-emergence within 3 days after planting.**

Apply before crop and weeds emerge. SPARTAN CHARGE herbicide may be applied with ground or aerial sprayers calibrated to deliver a minimum of 10 gallons finished spray per acre by ground application and 5 gallons of finished spray per acre by air. SPARTAN CHARGE herbicide may be applied with other pesticides registered for use on Spring Wheat.

**When applied according to directions, SPARTAN CHARGE herbicide will provide control of these key weeds:**

Russian Thistle

Kochia

For Information on other weeds not listed above, refer to Weed Controlled Section on the Section 3 label.

24(c) Registrant:



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000

## SPARTAN CHARGE Herbicide Use Rate Table (Spring Wheat Only) Post Plant Pre-emergence Applications

Broadcast Rate	SPARTAN CHARGE herbicide oz/A (lb ai/A)				
	Soil Texture				
% Organic Matter	Coarse	Medium (pH 7.0-7.6)	Medium (pH lower than 7.0)	Fine (pH 7.0 – 7.6)	Fine (pH lower than 7.0)
< 1.5 %	Do not use	Do not use	Do not use	Do not use	Do not use
1.5 - 3.0 %	Do not use	3.8 (0.10)	6.4 (0.17)	5.1 (0.14)	7.6 (0.21)
3.0+ %	Do not use	6.4 (0.17)	7.6 (0.21)	7.6 (0.21)	7.6 (0.21)

**Refer to the SPARTAN CHARGE herbicide section 3 label for information on soil types under the COARSE, MEDIUM, and FINE categories.**

### Restrictions:

- Applications must be made within 3 days after planting.
- Do not apply to Durum.
- Do not apply to soils classified as coarse. (Refer to section 3 label for information on soil types under the coarse, medium, and fine.)
- Do not apply to soils with a pH greater than 7.6
- Do not apply to soils with less than 1.5% organic matter.
- Do not apply more than 7.6 oz/A (0.21 lb ai/A) of SPARTAN CHARGE herbicide per crop year
- Do not apply SPARTAN CHARGE herbicide to Spring Wheat fields if SPARTAN CHARGE herbicide or additional sulfentrazone containing products have been previously applied within the same twelve-month period.
- Do not apply after crop emerges, or if the seedling is close to the soil surface.
- Do not incorporate to depths greater than 2 inches.
- Do not apply to frozen soils or to existing snow cover to prevent Spartan Charge herbicide runoff from rain or snow melt that may occur following application.
- Do not apply SPARTAN CHARGE herbicide after crop emergence, if the seedling is close to the soil surface, or the seed furrow is not closed properly as undesirable crop response may occur.
- Do not irrigate when Spring Wheat seedlings are close to emerging or cracking the soil surface

24(c) Registrant:



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000

## MANUFACTURER'S CONDITIONS AND RISK OF USE

NOTICE: This 24(C) label will only be utilized by growers located in the State of Idaho. SUCH GROWER USERS AGREE FMC CORPORATION WILL NOT BE HELD RESPONSIBLE FOR FAILURE TO PERFORM AND/OR CROP RESPONSE AND OR YIELD LOSSES RESULTING FROM THE USE OF SPARTAN CHARGE HERBICIDE AS A PRE-EMERGE TREATMENT TO SPRING WHEAT. Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability on the commercial SPARTAN CHARGE herbicide section 3 label and this 24c label before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product shall be followed carefully.

USE OF SPARTAN CHARGE HERBICIDE ON SPRING WHEAT IN ACCORDANCE WITH THIS LABELING MAY RESULT IN FAILURE TO PERFORM, CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SECTION 24(C) REGISTRATION BEFORE USING THIS PRODUCT ON THE CROP. FMC Corporation recommends that the user and/or grower test product in order to determine its suitability for such intended use.

FMC Corporation makes the Product available for use in the manner described in this Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of failure to perform, Crop injury, Crop yield reduction or Crop loss. The decision to use the Product in the manner described in the Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of failure to perform, Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this Labeling, you acknowledge and accept that, to the extent consistent with applicable law:

- (1) You assume all risk of failure to perform, Crop injury, Crop yield reduction and Crop loss;
- (2) FMC Corporation does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Labeling;
- (3) FMC Corporation does not make, and does not authorize any agent or representative to make, any warranties, expressed or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty;
- (4) FMC Corporation disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to failure to perform, Crop injury, Crop yield reductions and/or Crop loss, including without limitation incidental, consequential or special damages;
- (5) These Conditions and Risks of Use for 24(C) label registration supersede any contrary representations or recommendations by FMC Corporation, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Conditions and Risk of Use for this 24(C) Registration are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.

This product when used safflower may lead to crop injury, loss, or damage. FMC Corporation recommends that the user and/or grower test product in order to determine its suitability for such intended use. FMC Corporation makes the product available to the user and/or grower solely to the extent the benefit and utility, in sole opinion of the user and/or grower outweigh the extent of potential injury associated with the use of this product. The decision to use or not to use this herbicide must be made by each individual user and/or grower on the basis of possible crop injury from Spartan Charge herbicide,

24(c) Registrant:



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000

the severity of weed infestation, the cost of alternative weed controls and other factors, because of the risk of crop damage all such use is at the user's and/or grower's risk.

- Inadequate seed furrow closure or shallow planting (less than 1.5 inch) may result in undesirable crop response. As expected, poor growing conditions such as excessive moisture, low temperatures, soil compaction and diseases may also cause undesirable crop response.
- Adverse crop response may occur on coarse textured soils with low organic matter (less than 1.5%) and/or pH of 7.6 or higher, or on highly eroded soils, hilltops, or in areas of calcareous outcroppings. SPARTAN CHARGE herbicide use rates should be reduced in those areas or not applied in these areas at all. Inadequate seed furrow closure or shallow planting (less than 1.5 inch) may result in undesirable crop response and this product should not be applied. Poor growing conditions such as excessive moisture, low temperatures, soil compaction and diseases may also cause undesirable crop response.

These Crop Specific Use directions are based upon the interactive effects of SPARTAN CHARGE herbicide and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under General Application Instructions, Rotational Crop Guidelines, Replanting Instructions, Liability Disclaimer and any other section listed in the SPARTAN CHARGE herbicide section 3 label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with SPARTAN CHARGE herbicide. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on SPARTAN CHARGE herbicide under specific local conditions.

To the extent consistent with applicable law FMC or seller shall not be liable for any incidental, consequential, or special damages resulting from the use or handling of this product.

Label Code DR-4507 031921

24(c) Registrant:



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000