

Section 24(c) Special Local Need (SLN) Label

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

FIFRA 24(c) Special Local Need Label For Distribution and Use Only in the State of Idaho

Stinger[®] HL

EPA Reg. No. 62719-747

SLN ID-260006

Postemergence Broadleaf Weed Control in Radish Grown for Seed

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) label and the federal label for this product must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on the container label for EPA Reg. No. 62719-747.
- Follow all directions on this supplemental label.

Refer to the product label for additional Precautions and Restrictions, Mixing Instructions and Application Directions.

Active Ingredient:

clopypalid: 3,6-dichloro-2- pyridinecarboxylic acid, monoethanolamine salt.....	40.9%
Other Ingredients	59.1%
Total	100.0%

Acid Equivalent: clopypalid: 3,6-dichloro-2-pyridinecarboxylic acid - 31% (3 lb/gal)

Special Conditions and Risks of Use

USE OF Stinger[®] HL herbicide (THE “PRODUCT”) ON RADISH GROWN FOR SEED (THE “CROP”) MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.

Corteva Agriscience makes the Product available for use in the manner described in this Supplemental Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this Supplemental Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this Supplemental Labeling, you acknowledge and accept that:

- (1) You assume all risk of Crop injury, Crop yield reduction and/or Crop loss;
- (2) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Supplemental Labeling;

- 3) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaims all warranties, express or implied, including any implied warranties of fitness for a particular purpose or merchantability;
- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to inability to sell the Crop, Crop injury, Crop yield reduction and/or Crop loss;
- (5) These Special Conditions and Risks of Use supersede any contrary representations or recommendations by Corteva Agriscience, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use are required by Corteva Agriscience and not specified by the U.S. EPA or the State of Idaho.

Keep Out of Reach of Children

CAUTION

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Do not contaminate water used for irrigation or domestic purposes.

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow, or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

Directions for Use

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

Application Rates:

Target Weeds	Application Rate (pint/acre)	Use Restrictions
wild buckwheat chamomile clover	0.15 - 0.3 (0.09 - 0.19 lb ae/acre)	• Make 1 to 2 broadcast application per crop per year, not to exceed a total of 0.3 pint per acre

common cocklebur dandelion galinsoga prickly lettuce pineappleweed ragweed smartweed		(0.19 lb ae/acre) per year.
annual sowthistle ¹ Canada thistle ¹	0.2 – 0.3 (0.13 – 0.19 lb ae/acre)	

¹Suppression only.

Broadcast Application Rates: Apply uniformly with ground equipment in a minimum of 10 to 40 gallons of water per acre. For suppression of Canada thistle, apply after the majority of basal leaves have emerged but prior to bud stage and at least 30 days prior to harvest.

Application Timing: Risk of crop injury may increase when Stinger HL is applied during the bolting and/or flowering stages of crop growth. If sprays are needed at this time it is recommended to use a lower rate within the application rate range.

Specific Use Restrictions:

- Do not exceed 0.3 pint of Stinger HL per acre in a 12-month period.
- For use under this SLN label, do not tank mix Stinger HL with other herbicides.
- Do not apply Stinger HL within 6 to 8 hours of expected rainfall or irrigation.
- Chemigation: For use under this SLN label, do not apply this product through any type of irrigation system.
- For use under this SLN label, do not aerially apply Stinger HL.
- Avoid spray drift. Follow guidelines in the Avoiding Injury to Non-Target Plants section of the product label.
- Follow rotational crop restrictions and other use precautions and limitations on the product label for Stinger HL.
- Extreme growing conditions such as drought or near freezing temperatures prior to, at, and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth.
- Do not use plant residues, including hay or straw from treated areas, seed screenings, or manure or bedding straw from animals that have grazed or consumed forage from treated areas, for composting or mulching, where susceptible plants may be grown the following season. Do not spread manure from animals that have grazed or consumed seed screenings, forage or hay from treated areas on land used for growing susceptible broadleaf crops. To promote herbicidal decomposition, plant residues should be evenly incorporated or burned. Breakdown of clopyralid in crop residues or manure is more rapid under warm, moist soil conditions and may be enhanced by supplemental irrigation.
- Because crop injury may occur, do not use seed screenings/plant residue (even if composted) in growing media. Do not apply seed screenings/plant residue to land that might be planted to a susceptible crop.
- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

Special Crop Use Restrictions:

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a non-feed/non-food use. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have established pesticide residue tolerances for this crop. Consequently, no portion of this crop may be used or distributed for human consumption or animal feed. This restriction pertains to, but is not limited to: green chop, forage, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage, or seed screenings, and the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that screenings cannot be distributed or used for food or feed purposes. All restrictions listed in the Idaho Rules Governing Pesticide & Chemigation Use & Application, IDAPA 02.03.03.800 must be followed.

Any seed from a field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

“This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. **This seed, in whole, as sprouts, or in any form, may not be used for human consumption or animal feed.** Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration and other regulatory agencies.”

Corteva Agriscience LLC
9330 Zionsville Road
Indianapolis, Indiana 46268
1- 800-992-5994 (option #4)

™® Trademarks of Corteva Agriscience and its affiliated companies

R469-013
Approved:
Initial printing