



FIFRA Section 24(c) Special Local Need Label

Fontelis[®]

EPA Reg. No. 352-834

For Control of Foliar and Soil-borne Plant Diseases in Potatoes

For Distribution and Use Only in the State of Idaho.

This label is valid until December 31, 2026, or until otherwise amended, withdrawn, canceled or suspended.

24(c) Special Local Need Registration EPA SLN No. ID-220005

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label and the federal label for this product must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, and precautions on this supplemental label and the main EPA-registered label.

Environmental Hazards

This product is toxic to fish, aquatic invertebrates, and oysters. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff several weeks after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this chemical from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), and restricted-entry interval, and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes and socks
- Chemical resistant gloves (made of any waterproof material)

Directions for Use

Fontelis® is a broad-spectrum fungicide, recommended for control of foliar and soil-borne plant diseases in many crops, including potatoes, and has preventive, curative, and locally systemic activity. Fontelis must be applied in a regularly scheduled protective spray program in rotation with other fungicides

Refer to the product label for Precautions, Restrictions, Mixing Instructions, and Application Directions.

Disease Controlled or Suppressed	Product Rate fl oz/acre	Comments
Alternaria early blight and brown spot (Alternaria solani, Alternaria alternata) Gray mold (Botrytis cinerea) Powdery mildew (Erysiphe spp.)	10 to 24 fl oz	Begin applications prior to disease development and continue on a 7- to 14-day interval. Use higher rate and shorter interval when disease pressure is high.
Black dot (Colletotrichum coccodes)	14 to 24 fl oz	
Disease suppression White mold (Sclerotinia sclerotiorum)	14 to 24 fl oz	Make initial application at 100% full bloom of the primary inflorescence, or prior to row closure, and then again 14 days later.
Soil-borne diseases Rhizoctonia stem canker and black scurf (Rhizoctonia solani)	0.7 to 1.6 fl oz/1000 row-ft	At-plant, in-furrow application. Maximum rate per acre per application is 24 fl oz. For instructions, see "Soilborne/Seedling Disease Control for Tomatoes, Snap Beans and Bulb Onions" on the label affixed to the container for Fontelis and the "Soil Application Rates for Potatoes" below.

Make no more than 2 sequential applications of Fontelis before switching to a fungicide with a different mode of action. Minimum time (PHI) between application and harvest is 7 days. Do not exceed 72 fl oz (0.94 lb active ingredient) of Fontelis/acre per year from any combination of seed, soil, or foliar treatments.

Do not use Fontelis fungicide for an in-furrow application when penthiopyrad has been applied as a seed piece treatment for potatoes. Do not apply more than two in-season foliar potato applications of Fontelis at the maximum single application rate when penthiopyrad has been used as a seed piece treatment for potatoes. Do not apply more than 15 ounces of penthiopyrad active ingredient/acre/year to potatoes in total from any combination of seed, soil, or foliar treatments.

Soil Application Rates for Potatoes

Rate per 1000 row feet	Product per Acre (fl oz) ^a						
	22 " rows	30 " rows	32 " rows	34 " rows	36 " rows	38 " rows	40 " rows
0.7	16.7	12.2	11.4	10.9	10.2	9.6	9.2
1.2	28.6 ^b	20.9	19.6	18.5	17.4	16.5	15.7
1.6	-	27.9 ^c	26.1 ^d	24.6 ^e	23.0	22.0	21.1

^a Consult the maximum rate per acre allowed for the crop, and do not exceed that rate when using this application method.

^b In 22 inch rows, the highest rate for crops with 24 fl oz/acre maximums is 1.0 fl oz/1000 ft row, and for crops with 30 fl oz/acre maximums is 1.26 fl oz/1000 ft row.

^c In 30 inch rows, the highest rate for crops with 24 fl oz/acre maximums is 1.38 fl oz/1000 ft row.

^d In 32 inch rows, the highest rate for crops with 24 fl oz/acre maximums is 1.47 fl oz/1000 ft row.

^e In 34 inch rows, the highest rate for crops with 24 fl oz/acre maximums is 1.56 fl oz/1000 ft row.

Use Restrictions

- - Use this product only in commercial and farm plantings.
- - Do not use for home plantings.
- - Do not formulate this product into other end-use products.

Registrant

E. I. du Pont de Nemours and Company
 Chestnut Run Plaza
 974 Centre Road
 Wilmington, DE 19805

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

R613-014

Approved: __/__/__

Initial printing