

# FIFRA Section 24(c) Special Local Need (SLN) Label

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

For Distribution and Use Only in the State of Idaho

## Starane<sup>®</sup> Ultra

EPA Reg. No. 62719-577

EPA SLN No. ID-230001

### Postemergence Broadleaf Weed Control in Radish Grown for Seed, Including Daikon Grown for Seed

This label is valid until December 31, 2027 or until otherwise amended, withdrawn, cancelled, or suspended.

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) label and the federal label for this product must be in the possession of the user at the time of application.
- Read this SLN and the label affixed to the container for Starane<sup>®</sup> Ultra herbicide before applying.
- Follow all applicable directions, restrictions, and precautions on the container label for EPA Reg. No. 62719-577.
- Use of Starane Ultra according to this SLN label is subject to all use directions, restrictions, and precautions imposed by the label affixed to the container for Starane Ultra.

#### Active Ingredient(s):

fluroxypyr 1-methylheptyl ester: ((4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy)acetic acid, 1-methylheptyl ester..... 45.52%  
Other Ingredient(s) ..... 54.48%  
Total ..... 100.00%

Acid Equivalent: fluroxypyr: ((4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy)acetic acid – 31.59% - 2.8 lb/gal

#### Keep Out of Reach of Children

## WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

#### Environmental Hazards

This product is toxic to fish. Drift or runoff from treated areas may be hazardous to aquatic organisms and non-target plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below

the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

**DIRECTIONS FOR USE:**

**Application Rates:**

Target Weeds	Application Rate (pint/acre)	Use Restrictions
bindweed, field <sup>(1)</sup> bedstraw (cleavers) buckwheat, wild <sup>(1)</sup> morningglory mustard <sup>(1)</sup> nightshade species <sup>(1)</sup> pennycress, field <sup>(1)</sup> Russian thistle <sup>(1)</sup> velvetleaf	0.3-0.4 (0.11 - 0.14 lb ai/acre)	Make 1 to 2 broadcast application per crop per year, not to exceed a total of 0.7 pint per acre (0.24 lb ai/acre) per year.

<sup>1</sup>Suppression only.

**Broadcast Application Rates:** Apply in 3 or more gallons per acre by air or in 8 or more gallons per acre by ground equipment. Do not exceed 40 gallons per acre total spray volume. Use sufficient spray volume to provide thorough coverage and a uniform spray pattern. Inadequate spray volume and coverage may result in decreased weed control. As canopy and weed density increase, spray volume should be increased to obtain equivalent weed control. Use larger nozzle tips or decrease spraying speed to increase spray volume rather than increasing boom pressure. Refer to manufacturer's instructions for information on relationships between spray volume, and nozzle size and arrangement.

**Application Timing:** Apply to actively growing weeds. Extreme growing conditions such as drought or near freezing temperatures prior to, at and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. **Only weeds that are emerged at the time of application will be affected.** Foliage that is wet at the time of application may decrease control. Applications of Starane Ultra are rain-fast within 1 hour after application.

**Restrictions:**

- **Do not** exceed 0.7 pint per acre (0.24 lb ai/acre) per year.
- **Plant-back Restriction:** If replanting is required, plant only those crops listed on the product label for Starane Ultra within 120 days following application.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- **Preharvest Interval:** Do not apply closer than 14 days before harvest.
- Avoid spray drift. Follow guidelines in the "Precautions for Avoiding Spray Drift" section of the product label.
- Follow rotational crop restrictions and other use precautions and limitations on the product label for Starane Ultra.
- Extreme growing conditions such as drought or near freezing temperatures prior to, at, and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth.
- **Do not** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.
- No portion of the radish seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, leaves and seed screenings may be used or distributed for food or feed purposes.
- Seed resulting from plants/fields to which Starane Ultra was applied may not be distributed for human consumption or animal feed.

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training,

decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as natural rubber  $\geq$ 14 mil
- Shoes plus socks
- Protective eyewear

### **Special Conditions and Risks of Use**

**USE OF Starane® Ultra herbicide (THE “PRODUCT”) ON RADISH GROWN FOR SEED, INCLUDING DAIKON GROWN FOR SEED (THE “CROP”) MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.**

Corteva Agriscience makes the Product available for use in the manner described in this Section 24(c) or SLN labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this Section 24(c) or SLN labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this Labeling, you acknowledge and accept that:

- (1) You assume all risk of Crop injury, Crop yield reduction and/or Crop loss;
- (2) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Labeling;
- (3) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaims all warranties, express or implied, including any implied warranties of fitness for a particular purpose or merchantability;
- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to inability to sell the Crop, Crop injury, Crop yield reduction and/or Crop loss;
- (5) These Special Conditions and Risks of Use supersede any contrary representations or recommendations by Corteva Agriscience, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use are required by Corteva Agriscience and not specified by the U.S. EPA or the State of Idaho.

24(c) Registrant: Corteva Agriscience LLC  
9330 Zionsville Road  
Indianapolis, Indiana 46268  
800-992-5994 (option #4)

™® Trademarks of Corteva Agriscience and its affiliated companies

R357-056

Approved: \_\_/\_\_/\_\_

Initial Printing